arXiv:1509.02808v1 [math.AG] 9 Sep 2015

ON LOG K-STABILITY FOR ASYMPTOTICALLY LOG FANO VARIETIES

KENTO FUJITA

Abstract. The notion of asymptotically log Fano varieties was given by Cheltsov and Rubinstein. We show that, if an asymptotically log Fano variety (X, D) satisfies that D is irreducible and $-K_X - D$ is big, then X does not admit Kähler-Einstein edge metrics with angle $2\pi\beta$ along D for any sufficiently small positive rational number β. This gives an affirmative answer to a conjecture of Cheltsov and Rubinstein.

Contents

1. Introduction 1
2. Log K-stability 3
3. Constructing log semi test configurations 4
4. Examples 10

References 10

1. Introduction

The purpose of this article is to give a simple necessary criterion for log K-stability of $((X, D), -K_X-(1-\beta)D)$ with cone angle $2\pi\beta$ in the sense of [OS11], where X is projective log terminal and D is a reduced Weil divisor with $-K_X-(1-\beta)D$ ample. The motivation comes from a recent preprint of Cheltsov and Rubinstein [CR15], who treated the case that the dimension of X is equal to two. In this article, we show the following result.

Theorem 1.1 (=Theorem 3.9)**.** *Let X be a normal projective variety which is log terminal, D be a nonzero reduced Weil divisor on X which is $\mathbb{Q}$-Cartier, and $0 \leq \beta \leq 1$ be a rational number. Assume that the pair $(X, (1-\beta)D)$ is dlt, $-K_X-(1-\beta)D$ is ample, and $((X, D), -K_X-$*

Date: September 10, 2015.

2010 *Mathematics Subject Classification.* Primary 14J45; Secondary 14L24.

Key words and phrases. Fano varieties, K-stability, Kähler-Einstein metrics.

$(1-\beta)D)$ is log K-stable (resp. log K-semistable) with cone angle $2\pi\beta$. Then we have $\eta_\beta(D) > 0$ (resp. ≥ 0), where

$$\eta_\beta(D) := \beta\cdot\mathrm{vol}_X(-K_X-(1-\beta)D)-\int_0^\infty \mathrm{vol}_X(-K_X-(1-\beta+x)D)dx.$$

Note that vol_X *is the volume function (see* [Laz04]*).*

Theorem 1.1 immediately gives the following corollary.

Corollary 1.2 (see Corollary 3.11)**.** *Let X be a smooth projective variety and D be a nonzero reduced simple normal crossing divisor on X. Assume that $-K_X-(1-\beta)D$ is ample for any $0<\beta\ll 1$ and the divisor $-K_X-D$ is big. Then $((X,D),-K_X-(1-\beta)D)$ is not log K-semistable with cone angle $2\pi\beta$ for any $0<\beta\ll 1$ with $\beta\in\mathbb{Q}$. In particular, X does not admit Kähler-Einstein edge metrics with angle $2\pi\beta$ along D for any $0<\beta\ll 1$ with $\beta\in\mathbb{Q}$.*

Corollary 1.2 gives an affirmative answer for a conjecture of Cheltsov and Rubinstein for asymptotically log Fano varieties [CR13] with irreducible boundaries in any dimension. Although the following corollary is a special case of Corollary 1.2, we state the assertion for the readers' convenience.

Corollary 1.3 (see [CR13, Conjecture 1.11 (i)])**.** *Let (X,D) be an asymptotically log Fano variety with D irreducible, that is, X is a smooth projective variety and D is a smooth irreducible divisor on X such that $-K_X-(1-\beta)D$ is ample for any $0<\beta\ll 1$. If the divisor $-K_X-D$ is big, then X does not admit Kähler-Einstein edge metrics with angle $2\pi\beta$ along D for any $0<\beta\ll 1$ with $\beta\in\mathbb{Q}$.*

Remark 1.4. In [CR15, Theorem 1.6] (see also [CR15, Conjecture 1.5]), Cheltsov and Rubinstein proved Corollary 1.3 in dimension two by using a construction of flops on the deformation to the normal cone.

The strategy for the proof of Theorem 1.1 is essentially same as the strategy in [Fuj15]. We consider a kind of "log-version" of divisorial stability along D in the sense of [Fuj15]. We construct a specific log semi test configuration from certain section ring (see Remark 3.3) and calculate its log Donaldson-Futaki invariant explicitly by using the theory of "geography of models" (see Theorem 3.4).

Acknowledgments. The author thanks Professor Yanir Rubinstein who recommended to write down Section 4. The author acknowledges support from the Simons Center for Geometry and Physics, Stony Brook University at which some of the research for this paper was performed. The author is partially supported by a JSPS Fellowship for Young Scientists.

A *variety* stands for a reduced, irreducible, separated and of finite type scheme over the complex number field $\mathbb{C}$. For the theory of minimal model program, we refer the readers to [KM98]. For any Weil divisor E on a normal variety X, the *divisorial sheaf* on X is denoted by $\mathcal{O}_X(E)$. More precisely, the section $\Gamma(U, \mathcal{O}_X(E))$ on any open subscheme $U \subset X$ is given by the following:

$$\{f \in \mathbb{C}(X) \mid \operatorname{div}(f)|_U + E|_U \geq 0\},$$

where $\mathbb{C}(X)$ is the function field of X.

For varieties X_1 and X_2, let $p_i\colon X_1 \times X_2 \to X_i$ ($i = 1, 2$) be the projection morphisms.

2. Log K-stability

We recall the definition of log K-stability.

Definition 2.1 (see [OS11]). Let X be an n-dimensional normal projective variety, L be an ample line bundle on X, and D be a reduced Weil divisor on X.

(1) A coherent ideal sheaf $\mathcal{I} \subset \mathcal{O}_{X\times\mathbb{A}^1_t}$ is said to be a *flag ideal* if $\mathcal{I}$ is of the form

$$\mathcal{I} = I_M + I_{M-1}t^1 + \cdots + I_1 t^{M-1} + (t^M) \subset \mathcal{O}_{X\times\mathbb{A}^1_t},$$

where $I_M \subset \cdots \subset I_1 \subset \mathcal{O}_X$ is a sequence of coherent ideal sheaves of X.

(2) Let $m \in \mathbb{Z}_{>0}$, and let $\mathcal{I} \subset \mathcal{O}_{X\times\mathbb{A}^1}$ be a flag ideal. A *log semi test configuration* $((\mathcal{X}, \mathcal{D}), \mathcal{M})/\mathbb{A}^1$ *of* $((X, D), L^{\otimes m})$ *obtained by* $\mathcal{I}$ is given from the following data:
- $\Pi\colon \mathcal{X} \to X \times \mathbb{A}^1$ is the blowing up along $\mathcal{I}$, $\mathcal{D} \subset \mathcal{X}$ is given by the blowing up of $D \times \mathbb{A}^1$ along $\mathcal{I}|_{D\times\mathbb{A}^1}$, and $E \subset \mathcal{X}$ is the Cartier divisor defined by $\mathcal{O}_{\mathcal{X}}(-E) = \mathcal{I} \cdot \mathcal{O}_{\mathcal{X}}$,
- $\mathcal{M}$ is the line bundle on $\mathcal{X}$ defined by $\mathcal{M} := \Pi^* p_1^* L^{\otimes m} \otimes \mathcal{O}_{\mathcal{X}}(-E)$,

such that we require the following:
- $\mathcal{I}$ is not of the form (t^M), and
- $\mathcal{M}$ is semiample over $\mathbb{A}^1$.

(3) Assume that $((\mathcal{X}, \mathcal{D}), \mathcal{M})/\mathbb{A}^1$ is a log semi test configuration of $((X, D), L^{\otimes m})$ obtained by $\mathcal{I}$. Then the multiplicative group $\mathbb{G}_m$ naturally acts on $(\mathcal{X}, \mathcal{M})$ and $(\mathcal{D}, \mathcal{M}|_{\mathcal{D}})$. For $k \in \mathbb{Z}_{>0}$, let $w(k)$ be the total weight of $\mathbb{G}_m$-action on $H^0(\mathcal{X}_0, \mathcal{M}^{\otimes k}|_{\mathcal{X}_0})$ and $\tilde{w}(k)$ be the total weight of $\mathbb{G}_m$-action on $H^0(\mathcal{D}_0, \mathcal{M}^{\otimes k}|_{\mathcal{D}_0})$,

where $\mathcal{X}_0 \subset \mathcal{X}$ and $\mathcal{D}_0 \subset \mathcal{D}$ are the scheme-theoretic fibers at $0 \in \mathbb{A}^1$, respectively. It is known that, for $k \gg 0$, $w(k)$ (resp. $\tilde{w}(k)$) is a polynomial function of degree at most $n+1$ (resp. n). For $k \gg 0$, we set

$$\begin{aligned}\chi(X, L^{\otimes mk}) &= a_0k^n + a_1k^{n-1} + O(k^{n-2}),\\ \chi(D, L|_D^{\otimes mk}) &= \tilde{a}_0k^{n-1} + O(k^{n-2}),\\ w(k) &= b_0k^{n+1} + b_1k^n + O(k^{n-1}),\\ \tilde{w}(k) &= \tilde{b}_0k^n + +O(k^{n-1}).\end{aligned}$$

For any $\beta \in [0,1]$, we set the *log Donaldson-Futaki invariant* $\mathrm{DF}_\beta((\mathcal{X},\mathcal{D}),\mathcal{M})$ *with cone angle* $2\pi\beta$ as

$$\mathrm{DF}_\beta((\mathcal{X},\mathcal{D}),\mathcal{M}) := 2(b_0a_1 - b_1a_0) + (1-\beta)(a_0\tilde{b}_0 - b_0\tilde{a}_0).$$

(4) Let $\beta \in [0,1]$. $((X,D),L)$ is said to be *log K-stable* (resp. *log K-semistable*) *with cone angle* $2\pi\beta$ if $\mathrm{DF}_\beta((\mathcal{X},\mathcal{D}),\mathcal{M}) > 0$ (resp. ≥ 0) holds for any $m \in \mathbb{Z}_{>0}$, for any flag ideal $\mathcal{I}$, and for any log semi test configuration $((\mathcal{X},\mathcal{D}),\mathcal{M})/\mathbb{A}^1$ of $((X,D),L^{\otimes m})$ obtained by $\mathcal{I}$. For an ample $\mathbb{Q}$-divisor A on X, $((X,D),A)$ is said to be *log K-stable* (resp. *log K-semistable*) *with cone angle* $2\pi\beta$ if $((X,D),\mathcal{O}_X(aA))$ is so for some $a \in \mathbb{Z}_{>0}$ with aA Cartier (this definition does not depend on the choice of a).

The following theorem is important.

Theorem 2.2 (see [Ber12, CR15, OS11])**.** *Let X be a smooth projective variety, D be a reduced simple normal crossing divisor on X, and let $\beta \in [0,1] \cap \mathbb{Q}$. Assume that $-K_X - (1-\beta)D$ is ample and X admits Kähler-Einstein edge metrics with angle $2\pi\beta$ along D. Then $((X,D), -K_X - (1-\beta)D)$ is log K-semistable with cone angle $2\pi\beta$.*

3. Constructing log semi test configurations

In this section, from a pair (X,D), we construct a specific log semi test configuration via D. The construction is essentially in the same way as in [Fuj15, §3]. We fix the following condition:

Assumption 3.1. Let X be an n-dimensional normal projective variety which is log terminal, D is a nonzero reduced Weil divisor on X which is $\mathbb{Q}$-Cartier, and $\beta \in [0,1] \cap \mathbb{Q}$. Assume that the pair $(X,(1-\beta)D)$ is dlt and $-K_X - (1-\beta)D$ is ample.

Definition 3.2. Under Assumption 3.1, we set

$$\begin{aligned}\tau(D) &:= \sup\{\tau \in \mathbb{R}_{>0} \,|\, -K_X - \tau D \text{ big}\},\\ \tau_\beta(D) &:= \sup\{\tau \in \mathbb{R}_{>0} \,|\, -K_X - (1-\beta+\tau)D \text{ big}\}.\end{aligned}$$

It is obvious that $\tau_\beta(D) = \tau(D) - (1-\beta)$. We remark that $\tau(D) > 1$ holds if and only if the divisor $-K_X - D$ is big.

Remark 3.3. By [BCHM10, Corollary 1.1.9], the $\mathbb{C}$-algebra

$$\bigoplus_{\substack{k\in\mathbb{Z}_{\geq 0}\\ j\in\mathbb{Z}_{\geq 0}}} H^0(X, \mathcal{O}_X(\lfloor k(-K_X - (1-\beta)D) - jD\rfloor))$$

is finitely generated, where $\lfloor k(-K_X - (1-\beta)D) - jD\rfloor$ is the biggest $\mathbb{Z}$-divisor which is contained by $k(-K_X - (1-\beta)D) - jD$. We note that $H^0(X, \mathcal{O}_X(\lfloor k(-K_X - (1-\beta)D) - jD\rfloor)) = 0$ if $j > k\tau_\beta(D)$. Thus, there exists $r \in \mathbb{Z}_{>0}$ such that

- $L_\beta := r(-K_X - (1-\beta)D)$ is Cartier, and
- the $\mathbb{C}$-algebra

  $$\bigoplus_{\substack{k\in\mathbb{Z}_{\geq 0}\\ j\in[0, kr\tau_\beta(D)]\cap\mathbb{Z}}} H^0(X, \mathcal{O}_X(kL_\beta - jD))$$

  is generated by

  $$\bigoplus_{j\in[0, r\tau_\beta(D)]\cap\mathbb{Z}} H^0(X, \mathcal{O}_X(L_\beta - jD)).$$

From now on, we fix such r (and L_β).

Theorem 3.4 ([KKL12, Theorem 4.2])**.** *Under Assumption 3.1, there exist*

- *a sequence of rational numbers*

  $$0 = \tau_0 < \tau_1 < \cdots < \tau_m = \tau_\beta(D),$$

- *normal projective varieties $X_1, \ldots, X_m$ such that $X_1 = X$, and*
- *mutually distinct birational contraction maps $\phi_i\colon X \dashrightarrow X_i$ with $\phi_1 = \mathrm{id}_X$ $(1 \leq i \leq m)$*

such that the following hold:

- *for any $x \in [\tau_{i-1}, \tau_i]$, ϕ_i is a semiample model (see* [KKL12, Definition 2.3]*) of $-K_X - (1-\beta+x)D$, and*
- *if $x \in (\tau_{i-1}, \tau_i)$, then ϕ_i is the ample model (see* [KKL12, Definition 2.3]*) of $-K_X - (1-\beta+x)D$.*

Proof. By [BCHM10, Corollary 1.4.3], there exists a projective birational morphism $\sigma\colon \tilde{X} \to X$ such that σ is an isomorphism in codimension one and $\tilde{X}$ is $\mathbb{Q}$-factorial. Let $\tilde{D}$ be the strict transform of D on $\tilde{X}$. A semiample model (resp. the ample model) of $-K_{\tilde{X}} - (1-\beta+x)\tilde{D}$

is a semiample model (resp. the ample model) of $-K_X-(1-\beta+x)D$. Moreover, the $\mathbb{C}$-algebra

$$\bigoplus_{\substack{k\in\mathbb{Z}_{\geq 0}\\ j\in\mathbb{Z}_{\geq 0}}} H^0(X, \mathcal{O}_X(\lfloor k(-K_X-(1-\beta)D)-jD\rfloor))$$

is equal to the $\mathbb{C}$-algebra

$$\bigoplus_{\substack{k\in\mathbb{Z}_{\geq 0}\\ j\in\mathbb{Z}_{\geq 0}}} H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(\lfloor k(-K_{\tilde{X}}-(1-\beta)\tilde{D})-j\tilde{D}\rfloor)).$$

Thus we can apply [KKL12, Theorem 4.2]. □

We construct a log semi test configuration of $((X,D),\mathcal{O}_X(L_\beta))$ from D. For any $j\in[0,r\tau_\beta(D)]\cap\mathbb{Z}$, we set

$$I_j := \operatorname{Image}(H^0\left(X,\mathcal{O}_X(L_\beta-jD)\right)\otimes_{\mathbb{C}}\mathcal{O}_X(-L_\beta)\to\mathcal{O}_X),$$

where the homomorphism is the evaluation. Note that, for any $j\in[0,r\tau_\beta(D)]\cap\mathbb{Z}$, $I_j\subset\mathcal{O}_X(-jD)$ and

$$0\subset I_{r\tau_\beta(D)}\subset\cdots\subset I_1\subset I_0=\mathcal{O}_X$$

hold. For $k\in\mathbb{Z}_{>0}$ and $j\in[0,kr\tau_\beta(D)]\cap\mathbb{Z}$, we define

$$J_{(k,j)} := \sum_{\substack{j_1+\cdots+j_k=j,\\ j_1,\dots,j_k\in[0,r\tau_\beta(D)]\cap\mathbb{Z}}} I_{j_1}\cdots I_{j_k}\subset\mathcal{O}_X.$$

Lemma 3.5 (see [Fuj15, Lemma 3.3])**.** *The* $J_{(k,j)}\subset\mathcal{O}_X$ *is equal to*

$$\operatorname{Image}(H^0\left(X,\mathcal{O}_X(kL_\beta-jD)\right)\otimes_{\mathbb{C}}\mathcal{O}_X(-kL_\beta)\to\mathcal{O}_X).$$

In particular, we have

$$H^0\left(X,\mathcal{O}_X(kL_\beta-jD)\right)=H^0\left(X,\mathcal{O}_X(kL_\beta)\cdot J_{(k,j)}\right).$$

Proof. Set

$$V_{k,j} := H^0(X,\mathcal{O}_X(kL_\beta-jD))$$

for simplicity. We remark that, by the choice of $r\in\mathbb{Z}_{>0}$, the homomorphism

$$\bigoplus_{\substack{j_1+\cdots+j_k=j,\\ j_1,\dots,j_k\in[0,r\tau_\beta(D)]\cap\mathbb{Z}}} V_{1,j_1}\otimes_{\mathbb{C}}\cdots\otimes_{\mathbb{C}}V_{1,j_k}\to V_{k,j}$$

is surjective. For any $1\leq i\leq k$, the ideal sheaf I_{j_i} is nothing but

$$\operatorname{Image}(V_{1,j_i}\otimes_{\mathbb{C}}\mathcal{O}_X(-L_\beta)\to\mathcal{O}_X).$$

Thus the assertion follows. □

Set the flag ideal $\mathcal{I}$ such that

$$\mathcal{I} := I_{r\tau_\beta(D)} + I_{r\tau_\beta(D)-1}t^1 + \cdots + I_1 t^{r\tau_\beta(D)-1} + (t^{r\tau_\beta(D)}) \subset \mathcal{O}_{X\times\mathbb{A}^1_t}.$$

For any $k \in \mathbb{Z}_{>0}$, we have

$$\mathcal{I}^k = J_{(k,kr\tau_\beta(D))} + J_{(k,kr\tau_\beta(D)-1)}t^1 + \cdots + J_{(k,1)}t^{kr\tau_\beta(D)-1} + (t^{kr\tau_\beta(D)})$$

by the construction of $J_{(k,j)}$. Let $\Pi\colon \mathcal{X} \to X \times \mathbb{A}^1$ be the blowing up along $\mathcal{I}$ and let $E \subset \mathcal{X}$ be the Cartier divisor given by the equation $\mathcal{O}_{\mathcal{X}}(-E) = \mathcal{I}\cdot\mathcal{O}_{\mathcal{X}}$. Set $\mathcal{L}_\beta := \Pi^* p_1^*\mathcal{O}_X(L_\beta)\otimes\mathcal{O}_{\mathcal{X}}(-E)$. Let $\mathcal{D} \to D\times\mathbb{A}^1$ be the blowing up along $\mathcal{I}|_{D\times\mathbb{A}^1}$. We note that $\mathcal{I}|_{D\times\mathbb{A}^1} = (t^{r\tau_\beta(D)}) \subset \mathcal{O}_{D\times\mathbb{A}^1}$ since $I_j \subset \mathcal{O}_X(-jD) \subset \mathcal{O}_X(-D)$ for any $j > 0$. In particular, $\mathcal{D} \simeq D \times \mathbb{A}^1$ holds.

Lemma 3.6 (see [Fuj15, Lemma 3.4])**.** *$((\mathcal{X}, \mathcal{D}), \mathcal{L}_\beta)/\mathbb{A}^1$ is a log semi test configuration of $((X, D), L_\beta)$.*

Proof. Set $\alpha := p_2 \circ \Pi\colon \mathcal{X} \to \mathbb{A}^1$. It is enough to check that $\mathcal{L}_\beta$ is α-semiample. By Lemma 3.5, the homomorphism

$$H^0(X, \mathcal{O}_X(kL_\beta)\cdot J_{(k,j)}) \otimes_{\mathbb{C}} \mathcal{O}_X \to \mathcal{O}_X(kL_\beta)\cdot J_{(k,j)}$$

is surjective for any $k \in \mathbb{Z}_{>0}$ and $j \in [0, kr\tau_\beta(D)] \cap \mathbb{Z}$. Thus

$$H^0\left(X\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k\right) \otimes_{\mathbb{C}[t]} \mathcal{O}_{X\times\mathbb{A}^1} \to p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k$$

is surjective for any $k \in \mathbb{Z}_{>0}$. From [Laz04, Lemma 5.4.24], we have

$$\begin{aligned}
\alpha^*\alpha_*\mathcal{L}_\beta^{\otimes k} &\simeq \Pi^*(p_2)^*(p_2)_*(p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k)\\
&= \Pi^*\left(H^0\left(X\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k\right)\otimes_{\mathbb{C}[t]}\mathcal{O}_{X\times\mathbb{A}^1}\right)\\
&\twoheadrightarrow \Pi^*\left(p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k\right)\\
&\twoheadrightarrow \Pi^*p_1^*\mathcal{O}_X(kL_\beta)\otimes\mathcal{O}_{\mathcal{X}}(-kE) = \mathcal{L}_\beta^{\otimes k}
\end{aligned}$$

for $k \gg 0$. □

Definition 3.7. We say the log semi test configuration $((\mathcal{X}, \mathcal{D}), \mathcal{L}_\beta)/\mathbb{A}^1$ the *basic log semi test configuration of* $((X, D), \mathcal{O}_X(L_\beta))$ *via* D.

Now we calculate the log Donaldson-Futaki invariant of the basic log semi test configuration $((\mathcal{X}, \mathcal{D}), \mathcal{L}_\beta)/\mathbb{A}^1$ of $((X, D), \mathcal{O}_X(L_\beta))$ via D. By the asymptotic Riemann-Roch theorem, we have

$$a_0 = \frac{(L_\beta^{\cdot n})}{n!}, \quad a_1 = \frac{(L_\beta^{\cdot n-1}\cdot -K_X)}{2\cdot(n-1)!}, \quad \tilde{a}_0 = \frac{(L_\beta^{\cdot n-1}\cdot D)}{(n-1)!}.$$

(We follow the notation in Definition 2.1.) By [Odk13, §3],

$$\begin{aligned}
w(k) &= -\dim\left(\frac{H^0(X\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta))}{H^0(X\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta)\cdot\mathcal{I}^k)}\right)\\
&= -kr\tau_\beta(D)\cdot h^0(X, \mathcal{O}_X(kL_\beta)) + v(k),
\end{aligned}$$

where

$$v(k) := \sum_{j=1}^{kr\tau_\beta(D)} h^0(X, \mathcal{O}_X(kL_\beta - jD)).$$

By the same argument,

$$\begin{aligned} \tilde{w}(k) &= -\dim\left(\frac{H^0(D\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta)|_D)}{H^0(D\times\mathbb{A}^1, p_1^*\mathcal{O}_X(kL_\beta)|_D\cdot(t^{kr\tau_\beta(D)}))}\right)\\ &= -kr\tau_\beta(D)\cdot h^0(D, \mathcal{O}_X(kL_\beta)|_D). \end{aligned}$$

Thus

$$\tilde{b}_0 = -r\tau_\beta(D)\frac{(L_\beta^{\cdot n-1}\cdot D)}{(n-1)!}.$$

We set $v(k) = v_0k^{n+1} + v_1k^n + O(k^{n-1})$. We calculate the values v_0 and v_1. Let $L_{\beta,i}$ and D_i be the divisors on X_i which are the push-forwards of L_β and D, respectively. For $k \gg 0$ sufficiently divisible, by [KKL12, Remark 2.4 (i)] and [Fuj15, Proposition 4.1], $v(k)$ is equal to

$$\begin{aligned} &\sum_{i=1}^{m}\sum_{j=kr\tau_{i-1}+1}^{kr\tau_i} h^0\left(X_i, \mathcal{O}_{X_i}\left(kL_{\beta,i} - jD_i\right)\right)\\ = &\sum_{i=1}^{m}\Big(\frac{(kr)^{n+1}}{n!}\int_{\tau_{i-1}}^{\tau_i}\left(((1/r)L_{\beta,i} - xD_i)^{\cdot n}\right)dx\\ &-\frac{(kr)^n}{2\cdot(n-1)!}\int_{\tau_{i-1}}^{\tau_i}\left(((1/r)L_{\beta,i} - xD_i)^{\cdot n-1}\cdot(K_{X_i}+D_i)\right)dx\Big)\\ &+O(k^{n-1}). \end{aligned}$$

This implies that

$$\begin{aligned} v_0 &= \frac{r^{n+1}}{n!}\sum_{i=1}^{m}\int_{\tau_{i-1}}^{\tau_i}\left(((1/r)L_{\beta,i} - xD_i)^{\cdot n}\right)dx,\\ v_1 &= \frac{-r^n}{2\cdot(n-1)!}\sum_{i=1}^{m}\int_{\tau_{i-1}}^{\tau_i}\left(((1/r)L_{\beta,i} - xD_i)^{\cdot n-1}\cdot(K_{X_i}+D_i)\right)dx. \end{aligned}$$

Thus we have

$$\begin{aligned} &\mathrm{DF}_\beta((\mathcal{X},\mathcal{D}),\mathcal{L}_\beta)\\ = &\ 2(v_0a_1 - v_1a_0) + (1-\beta)(a_0\tilde{b}_0 - (v_0 - r\tau_\beta(D)a_0)\tilde{a}_0)\\ = &\ \frac{n\cdot r^n(L_\beta^{\cdot n})}{(n!)^2}\sum_{i=1}^{m}\int_{\tau_{i-1}}^{\tau_i}(\beta - x)\left((-K_{X_i} - (1-\beta+x)D_i)^{\cdot n-1}\cdot D_i\right)dx. \end{aligned}$$

Lemma 3.8 (cf. [Fuj15, Theorem 5.2])**.** *We have*

$$\eta_\beta(D) = n\sum_{i=1}^{m}\int_{\tau_{i-1}}^{\tau_i}(\beta - x)\left((-K_{X_i} - (1-\beta+x)D_i)^{\cdot n-1}\cdot D_i\right)dx.$$

Proof. By [KKL12, Remark 2.4 (i)], we have

$$\mathrm{vol}_X(-K_X - (1-\beta+x)D) = ((-K_{X_i} - (1-\beta+x)D_i)^{\cdot n})$$

for any $x \in [\tau_{i-1}, \tau_i]$. From partial integration, we have

$$\begin{aligned} & n\sum_{i=1}^{m}\int_{\tau_{i-1}}^{\tau_i}(\beta - x)\left((-K_{X_i} - (1-\beta+x)D_i)^{\cdot n-1}\cdot D_i\right)dx \\ = & \sum_{i=1}^{m}\Big([(x-\beta)\,\mathrm{vol}_X(-K_X-(1-\beta+x)D)]_{\tau_{i-1}}^{\tau_i} \\ & -\int_{\tau_{i-1}}^{\tau_i}\mathrm{vol}_X(-K_X-(1-\beta+x)D)dx\Big) = \eta_\beta(D). \end{aligned}$$

We remark that $\mathrm{vol}_X(-K_X - (1-\beta+x)D) = 0$ if $x \geq \tau_\beta(D)$. □

Therefore we have obtained the following.

Theorem 3.9. *Under Assumption 3.1, assume that* $((X, D), -K_X - (1-\beta)D)$ *is log K-stable (resp. log K-semistable) with cone angle* $2\pi\beta$*. Then* $\eta_\beta(D) > 0$ *(resp.* ≥ 0*) holds.*

Remark 3.10. If $\beta = 1$, then the value $\eta_1(D)$ is noting but the value $\eta(D)$ in [Fuj15, Definition 1.1].

Corollary 3.11. *Let* X *be a normal projective variety which is log terminal,* D *is a nonzero reduced Weil divisor on* X *which is* $\mathbb{Q}$*-Cartier. Assume that* $(X, (1-\beta)D)$ *is klt and* $-K_X - (1-\beta)D$ *is ample for any* $0 < \beta \ll 1$*. Moreover, we assume that* $-K_X - D$ *is big. Then for any* $0 < \beta \ll 1$ *rational number,* $((X, D), -K_X - (1-\beta)D)$ *is not log K-semistable with cone angle* $2\pi\beta$*.*

Proof. We have $\eta_\beta(D) = \eta_+(\beta) - \eta_-$, where

$$\begin{aligned} \eta_+(\beta) &:= \beta\cdot\mathrm{vol}_X(-K_X-(1-\beta)D) - \int_{1-\beta}^{1}\mathrm{vol}_X(-K_X - xD)dx, \\ \eta_- &:= \int_1^\infty \mathrm{vol}_X(-K_X - xD)dx. \end{aligned}$$

If $-K_X - D$ is big, then $\eta_- > 0$. On the other hand, $\lim_{\beta\to 0}\eta_+(\beta)$ is equal to zero. Thus the assertion follows from Theorem 3.9. □

Corollary 1.2 is immediately obtained from Theorem 2.2 and Corollary 3.11.

4. Examples

We see some examples.

Example 4.1. Let X be an n-dimensional Fano manifold, and let D be a smooth divisor on X with $-K_X \sim_{\mathbb{Q}} lD$ for some $l \in [1, n+1] \cap \mathbb{Q}$. Then $-K_X - (1-\beta)D$ is ample for any $\beta \in (0,1]$. In this case, we have

$$\eta_\beta(D) = \frac{n}{n+1} \operatorname{vol}_X(-K_X - (1-\beta)D)\left(\beta - \frac{l-1}{n}\right).$$

If $\beta < (l-1)/n$, then $\eta_\beta(D) < 0$ holds.

Remark 4.2. In Example 4.1, if $D \sim -K_X$ (i.e., $l = 1$), then $\eta_\beta(D) > 0$ for any $\beta \in (0,1]$. Thus our argument does not give any destabilizing information in this case. Compare with [CR15, §6].

Example 4.3. Let $Y := \mathbb{P}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(1))$, C be a section of the $\mathbb{P}^1$-bundle $Y/\mathbb{P}^1$ with $(C^{\cdot 2}) = 1$, $\pi \colon X \to Y$ be the blowing up along $p \in Y$ with $p \in C$, E be the exceptional divisor of π, and set $D := \pi_*^{-1}C$. Then $-K_X - (1-\beta)D$ is ample for any $\beta \in (0,1]$. (The pair (X, D) is nothing but [CR13, (I8B.1)].) In this case, $\tau_1 = \beta$, $X_2 = Y$, $\tau_2 = \tau_\beta(D) = 1 + \beta$ and

$$\operatorname{vol}_X(-K_X - xD) = \begin{cases} -4x + 7 & \text{if } x \in [0,1], \\ x^2 - 6x + 8 & \text{if } x \in [1,2]. \end{cases}$$

Thus $\eta_\beta(D) = 2(\beta^2 - 2/3)$. For example, if we set $\beta := 1/2$, then $\eta_{1/2}(D) < 0$ holds. In this case ($\beta = 1/2$), we can check that $r := 2$ satisfies the condition in Remark 3.3 and the corresponding flag ideal $\mathcal{I}$ is of the form

$$\mathcal{I} = \mathcal{O}_X(-3D - 2E) + \mathcal{O}_X(-2D - E)t^1 + \mathcal{O}_X(-D)t^2 + (t^3).$$

References

[BCHM10] C. Birkar, P. Cascini, C. D. Hacon and J. M$^{\mathrm{c}}$Kernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468.

[Ber12] R. Berman, *K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics*, arXiv:1205.6214; to appear in Invent. Math.

[CR13] I. A. Cheltsov and Y. A. Rubinstein, *Asymptotically log Fano varieties*, arXiv:1308.2503; to appear in Adv. Math.

[CR15] I. A. Cheltsov and Y. A. Rubinstein, *On flops and canonical metrics*, arXiv:1508.04634.

[Fuj15] K. Fujita, *On K-stability and the volume functions of* $\mathbb{Q}$*-Fano varieties*, arXiv:1508.04052.

[KKL12] A.-S. Kaloghiros, A. Küronya and V. Lazić, *Finite generation and geography of models*, arXiv:1202.1164; to appear in Advanced Studies in Pure Mathematics, Mathematical Society of Japan, Tokyo.

[KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math, vol.134, Cambridge University Press, Cambridge, 1998.

[Laz04] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, Ergebnisse der Mathematik und ihrer Grenzgebiete. (3) **48**, Springer, Berlin, 2004.

[Odk13] Y. Odaka, *A generalization of the Ross-Thomas slope theory*, Osaka. J. Math. **50** (2013), no. 1, 171–185.

[OS11] Y. Odaka and S. Sun, *Testing log K-stability by blowing up formalism*, arXiv:1112.1353; to appear in Ann. Fac. Sci. Toulouse Math.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN
E-mail address: `fujita@math.kyoto-u.ac.jp`